

Prostate Cancer Discussion Guide

PSA Level: _____ Date: _____

Testosterone Level: _____ Date: _____

Has my cancer spread outside the prostate?

- Yes (metastatic) No (non-metastatic)

Is my cancer still responding to initial hormone therapy?

- Yes (castration-sensitive) No (castration-resistant)

What treatment options are available for my grade/stage of cancer?

PROVENGE

Indication: PROVENGE® (sipuleucel-T) is a prescription medicine used to treat certain men with advanced prostate cancer. PROVENGE is an established cellular immunotherapy and is customized to each individual by using his own immune cells.

How is PROVENGE different than other treatments I may be taking for my prostate cancer?

Am I eligible for PROVENGE? Yes No Why or why not? _____

Criteria:

- Have disease progression on hormone therapy (ADT)
- Have testosterone levels that are less than 50 ng/dL
- Diagnosed with metastatic disease
- Not taking drugs for cancer-related pain

How do I get started on PROVENGE? _____

Notes: _____

What am I currently doing to monitor my prostate cancer?

- | | |
|---|------------------|
| <input type="checkbox"/> PSA Test | Frequency: _____ |
| <input type="checkbox"/> Testosterone Test | Frequency: _____ |
| <input type="checkbox"/> Conventional Imaging | Frequency: _____ |
| <input type="checkbox"/> Bone Scan | Frequency: _____ |
| <input type="checkbox"/> CT Scan | Frequency: _____ |
| <input type="checkbox"/> Next Generation Imaging (PET Scan with Diagnostic Imaging Agent) | Frequency: _____ |
| <input type="checkbox"/> Other: _____ | Frequency: _____ |
| <input type="checkbox"/> Other: _____ | Frequency: _____ |

Helpful Tip: Set reminders in your calendar for the necessary tests

PROVENGE[®]
(sipuleucel-T)

PROVENGE®

(sipuleucel-T)

INDICATION

PROVENGE® (sipuleucel-T) is a prescription medicine used to treat certain men with advanced prostate cancer. PROVENGE is an established cellular immunotherapy and is customized to each individual by using his own immune cells.

IMPORTANT SAFETY INFORMATION

Before receiving PROVENGE®, tell your doctor about any medical conditions, including heart or lung problems, or if you have had a stroke.

Tell your doctor about any medicines you take, including prescription and nonprescription drugs, vitamins, or dietary supplements.

The most common side effects of PROVENGE include chills, fatigue, fever, back pain, nausea, joint ache, and headache. These are not all the possible side effects of PROVENGE treatment.

PROVENGE is made from your own immune cells, which are collected during a process called leukapheresis. The cells are processed, returned, and then infused back into the patient through an IV (intravenous) infusion about 3 days later. This process is completed in 3 cycles, about 2 weeks apart. Each infusion takes approximately 1 hour and requires 30 minutes of post-infusion monitoring.

PROVENGE infusion can cause serious reactions. Tell your doctor right away if you:

- Have signs of a heart attack or lung problems, such as trouble breathing, chest pains, racing or irregular heart beats, high or low blood pressure, dizziness, fainting, nausea, or vomiting
- Have signs of a stroke, such as numbness or weakness on one side of the body, decreased vision in one eye, or difficulty speaking
- Develop symptoms of thrombosis which may include: pain and/or swelling of an arm or leg with warmth over the affected area, discoloration of an arm or leg, shortness of breath, chest pain that worsens or deep breathing
- Have signs of infection such as a fever over 100°F, redness or pain at the infusion or collection sites

Tell your doctor about any side effect(s) that concerns you or does not go away. For more information, talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



Scan here for full
Prescribing Information
or visit
www.PROVENGE.com/pi

Dendreon®